

Asian Journal of Research in Chemistry and Pharmaceutical Sciences

Journal home page: www.ajrcps.com

<https://doi.org/10.36673/AJRCPS.2021.v09.i03.A13>



STANNOUS CHLORIDE INDUCES ALTERATIONS IN BODY WEIGHT AND HEMATOLOGICAL PARAMETERS IN MALE RABBIT

Mariam A. Ibrahim^{*1}, Eman. K. Saad¹, Fayrouz A. Khaled¹

^{1*}Department of Chemistry, Faculty of Science, Omar Al-Mokhtar University, El -Beida-Libya.

ABSTRACT

Stannous chloride (SnCl_2) is widely used in daily human life, for example, to conserve soft drink, in food manufacturing and biocidal preparations. Animal in this study (ten white rabbits) were divided at random into two equal groups (each group five rabbits). The first group designed to be considered as a control. The second group was created to study the effect of stannous chloride (20mg/kg body weight) for 12 weeks. Overall means indicated that treatment with stannous chloride caused significant ($P < 0.05$) decrease in body weight and relative weight of liver, kidney, spleen, testes, and heart compared to control animals. Additionally, this study indicated that treatment with stannous chloride (SnCl_2) caused significant increase in white blood cells (WBCs) and insignificant decrease in haemoglobin (Hb), red blood cells (RBCs), packed cell volume (PCV) and platelets. The study confirmed the detrimental effects of stannous chloride on the body weight, relative organ weight and hematological parameters on male rabbits.

KEYWORDS

Stannous chloride, Rabbits, Relative organ weight and Hematological parameters.

Author for Correspondence:

Mariam A. Ibrahim,
Department of Chemistry,
Faculty of Science, Omar Al-Mokhtar University,
El -Beida-Libya.

Email: mariam.ibrahim@omu.edu.ly

INTRODUCTION

Tin derivatives and compounds are shown an important role in life activity. For long time it use in plated containers and alloys¹. Many of the inorganic tin compounds are enter in industrial processes for the strengthening of glass, as a base for colors, as catalysts, as stabilizers in perfumes and soap and as dental cryogenic agents. One of the common stannous ions, is the stannous chloride (SnCl_2) salt form, is extensively used as a reducing agent to label radiotracers with technetium-99m (^{99m}Tc). These radiotracers can be used as radio pharmaceuticals in nuclear medicine procedures. In

this case, absorption of this complex was confirmed, because it is intravenously administered in humans, although biological effects of these agents have not been fully understood².

The chemical structure of SnCl₂ has been identified as a compound that has a lone pair of electrons. In the solid state, crystalline SnCl₂ forms chains linked via chloride bridges as shown in Figure No.1³.

Tin(II) chloride, also known as stannous chloride, is a white crystalline solid with the formula SnCl₂. In addition, using stannous chloride induced a dose-dependent increase in SnCl₂ as a reducing agent (in acid solution) is widespread, and in electrolytic baths for tin-plating. SnCl₂ should not be confused with the other chloride of tin; tin(IV) chloride or stannic chloride (SnCl₄). In the last 25 years, considerable efforts have been made to understand the effect of tin, tin compounds and Sn²⁺ on human cells⁴ and in the environment the food industry uses stannous chloride as a preservative (e.g., in soft drinks) and in some fluoride toothpastes⁵. Possible joint consumption of these compounds, therefore, makes relevant information on their effect, isolated or in interaction, even with other organisms used as indicators⁶. The nano tin compounds effect on human health and the environment are under investigation due to the complexity of measuring the biological effect of nanoparticles/nanomaterials. In the environment, the size and state of agglomeration/aggregation vary, affecting the bioavailability of the nanoparticles and their access and accumulation in cells or their release from the body⁷. Stannous is used for the treatment of rheumatoid arthritis and other inflammatory diseases⁸. With the use of stannous ointment, the local metal exposure can reach 0.15g day⁻¹. Humans are exposed to stannous mostly through tin-lined cans but also through toothpaste, perfumes and food additives or to organotin compounds via plastic pipes, paints and pesticides. Stannous overload can cause anemia as well as liver and kidney problems⁹. Moreover, stannous chloride has an effect on inducing a dose-dependent increase in the MN frequency in peripheral erythrocytes of adult zebra fish (*Danio rerio*) after 120 exposures to doses $\geq 50\text{M}^{10}$. The use of herbal products is

gaining popularity around the world, as they are considered to be effective and to have few side-effects¹¹. Variation in blood constituents gives a rapid and good picture reflecting the physiological status of the organism because these changes develop more quickly in response to toxicants than any apparent morphological changes¹². Janssen *et al* investigated the effect of stannous chloride at 0,250,500ppm Sn²⁺. They found that haemoglobin decreased significantly¹³. Rader and his group, studied the anti-nutritive effect of dietary tin. He reported that a variety of effects of tin including variations in the activity of heme oxygenase and δ -aminolevulinic acid dehydratase and also tin has adverse effects on metabolism of essential trace minerals including iron¹⁴.

MATERIAL AND METHODS

In this study stannous chloride (SnCl₂) was used. SnCl₂ (purity 400g/L) was brought from chemistry department, faculty of science, Omar Al-Mokhtar University, El-Beida-Libya. Ten Male New Zealand adult White rabbits age of 6 months and weighing between (1.891 \pm 27.6Kg) were included in the study. All rabbits were housed individually in cages and checked the weight every week throughout 3-months experimental period. The first group was used as control, while, groups 2 were treated with SnCl₂ by gavage at a dose of 20mg/kg B.W/day (1/50 of SnCl₂) lethal dose¹⁵. At the end of the experimental period body weight of rabbits were recorded. Animals were sacrificed by decapitation and liver, lung, kidney, testes and brain were immediately removed and weighed then the organs weight ratio was calculated. All relative weight of organs (%) were calculated as g/100g body weight. Two tubes were used to collect the blood samples as follows: one containing EDTA (anti-coagulant) and the other containing Heparin (anti-coagulant). No coagulated blood by EDTA was tested shortly after collection by Particle counter (from ERMA INC.-Tokyo. Model PCE-210) for measuring total leukocyte counts (TLC), total erythrocyte count (TEC), platelet count (PLT), haemoglobin (Hb), packed cells volume (PCV). Statistical analysis. Where viable, statistical analysis was carried out in

Minitab software; statistical significance was determined by using one way ANOVA analysis.

RESULTS AND DISCUSSION

Effect of stannous chloride (SnCl₂) on body weight and relative organ weights

The changes in body weight (BW) and the relative weights of liver, kidney, lung, spleen, testes, and heart of male rabbits. The relative organ weights (%) were calculated as g/100g body weight throughout the 12week experimental period of rabbits treated with Stannous chloride (SnCl₂) were summarized in (Table No.1). Overall means indicated that treatment with SnCl₂ caused significant (P<0.05) decrease in BW and relative weight of liver, kidney, spleen, testes, and heart compared to control animals.

Effect of ginseng, stannous chloride and their combination on hematological parameters

Table No.2 presents the hematological parameters of male rabbits treated with ginseng, SnCl₂ Results indicated that treatment with stannous chloride (SnCl₂) caused significant increase in white blood cells (WBCs) and insignificant decrease in haemoglobin (Hb), red blood cells (RBCs), packed cell volume (PCV) and platelets. Similar effects were observed with values of MCV and MCH in SnCl₂. While values of MCHC was not showed any significant deference.

Discussion

In this study, was confirmed the treatment with SnCl₂ caused significant reductions in body weight (BW) and relative organs weight (ROW) (Table No.1). The reduction in BW and ROW of the SnCl₂ treated rabbits is in agreement with the finding of¹⁶. Relative organs weight (ROW) were reduced by SnCl₂ treatment. Similar results were obtained by Beynen *et al*, Yu and Beynen and Omura *et al*, in rabbits¹⁷⁻¹⁹. Also, testicular degeneration was observed in rats receiving 10mg of tin(II) chloride per kg in the feed for 12 weeks²⁰.

The present study showed that SnCl₂ caused decrease RBCs, Hb, PCV and PLT (Table No.2) agreement with Beynen *et al*, who found that iron status (tissue iron, haemoglobin, hematocrit, red blood cell count, plasma iron, total iron binding capacity and transferrin saturation) in rabbits was not influenced by dietary tin concentrations < 100mg Sn/kg diet as SnCl₂ for 28 days. Higher dietary intake of tin caused a decrease in these parameters¹⁷. Food intake and body weights were not reported. A study in waster rabbits fed on diets containing various concentrations of tin (1, 10, 50, 100 and 200mg Sn/kg as SnCl₂) for 28 days showed that iron, copper and Zink tissue and plasma concentrations were seemingly unaffected at 1mg and slightly decreased at 10mg Sn/kg diet (~ 0.7mg Sn/kg body weight/day). Greater effects were reprted at 50mg/kg diet (~ 3.5mg Sn/kg body weight/day). The blood haemoglobin concentration and percentage transferrin saturation decreased in a linear manner as the level of dietary Sn increased. Janssen *et al*, Investigated the effects of 0, 250 or 500mg Sn/kg diet (as SnCl₂) in a 4-week study on weanling Wistar rabbits, Haemoglobin was decreased and body weights reduced in a dose-related way in the tin-fed groups. Crypt depth, villus length and cell turnover were increased in parts of the intestine. In week 4, the estimated doses of tin were about 25 and 50mg Sn/kg body weight/day, respectively¹³.

Table No.1: Body weight (BW) and relative weight of kidney, liver, lung, brain, heart, testes and spleen of male rabbits treated with stannous chloride (SnCl₂)

S.No	Parameter	Control	SnCl ₂
1	BW (gm)	1.891± 35.64 ^a	1.756 ± 58.65 ^a
2	Kidney (g/100gm)	11.84± 1.275 ^a	10.34± 0.382 ^a
3	Liver (g/100gm)	42.78 ± 0.697 ^a	39.10 ± 1.453 ^a
4	Lung (g/100gm)	8.520 ± 0.736 ^{ab}	7.600 ± 0.510 ^b
5	Heart (g/100gm)	6.400 ± 0.872 ^a	5.900 ± 0.332 ^a
6	Testes (g/100gm)	4.080 ± 0.972 ^a	3.604 ± 0.713 ^a
7	Spleen (g/100gm)	4.080 ± 0.972 ^a	3.604 ± 0.713 ^a

Values are expressed as means ± SE; n = 5 for each treatment group. Mean values within a row not sharing a common superscript letters (a, b) were significantly different, p<0.05.

Table No.2: Changes Complete blood counts Red blood cells (RBC), white blood cells (WBC), packed cell volume (PCV), platelets count (PLT), hemoglobin (Hb), of male rabbits treated with stannous chloride (SnCl₂)

S.No	Parameters	Control	SnCl ₂
1	RBC ×10 ⁶ (μl)	5.58± 0.374 ^{ab}	4.55±0.096 ^c
2	Hb (g/dl)	13.27±0.20 ^{ab}	12.08±0.31 ^c
3	PCV×10 ³ (μl)	40.64±0.252 ^a	36.70±0.844 ^b
4	WBC ×10 ³ (μl)	8.63± 0.47 ^b	7.47± 0.27 ^c
5	PLT ×10 ³ (μl)	273.2±9.35 ^a	177.6±7.45 ^b
6	MCV (fl)	75.44±2.35 ^{ab}	81.18±1.91 ^a
7	MCH (pg)	24.45±2.23 ^b	29.46±0.42 ^a
8	MCHC (dl)	32.69± 0.45 ^a	32.82±0.57 ^a

Values are expressed as means ± SE; n = 5 for each treatment group. Mean values within a row not sharing a common superscript letters (a, b) were significantly different, p<0.05.




Figure No.1: Chemical structure of stannous chloride³

CONCLUSION

Our finding confirmed a marked effect of SnCl₂ on decreasing a body weight and relative weight of some organs and treatment with SnCl₂ cause an obvious changes in all hematological parameters by increasing white blood cells(WBCs) and decreasing in hemoglobin (Hb), red blood cells (RBCs), packed cell volume (PCV) and platelets.

ACKNOWLEDGEMENT

The authors wish to express their sincere gratitude to Department of Chemistry, Faculty of Science, Omar Al-Mokhtar University, El -Beida-Libya for providing necessary facilities to carry out this research work.

CONFLICT OF INTEREST

We declare that we have no conflict of interest.

BIBLIOGRAPHY

1. El-Demerdash F M, Yousef M I, Zoheir M A. Stannous chloride induces alterations in enzyme activities, lipid peroxidation and histopathology in male rabbit: Antioxidant role of Vitamin C, *Food Chem. Toxicol*, 43(12), 2005, 1743-1752.
2. Assis M L B, De Mattos J C P, Caceres M R, Dantas F J S, Asad L M B O, Asad N R, Bezerra R J A C, Caldeira-De-Araujo A, Bernardo-Filho M. Adaptive response to h₂o₂ protects against SnCl₂ damage: The OxyR system involvement, *Biochimie*, 84(4), 2002, 291-294.
3. Leger J M, Haines J, Atouf A. The high pressure behaviour of the cotunnite and post-cotunnite phases of PbCl₂ and SnCl₂, *J. Phys. Chem. Solids*, 57(1), 1996, 7-16.
4. Reynolds A S, Pierre T H, McCall R, Wu J, Gato W E. Evaluating the cytotoxicity of tin dioxide nanofibers, *J. Environ. Sci. Heal. - Part A Toxic/Hazardous Subst. Environ. Eng*, 53(11), 2018, 986-991.
5. Gallo A. Toxicity of marine pollutants on the ascidian oocyte physiology: An electrophysiological approach, *Zygote*, 26(1), 2018, 14-23.
6. Benowitz N L. Clinical pharmacology of caffeine, *Annual Review of Medicine*, 41, 1990, 277-288.
7. Ribeiro A R, Leite P E, Falagan-Lotsch P, Benetti F, Micheletti C, Budtz H C, Jacobsen N R, Lisboa-Filho P N, Rocha L A, Kuhnel D *et al.* Challenges on the toxicological predictions of engineered nanoparticles, *NanoImpact*, 8, 2017, 59-72.
8. Heusser P, Scheffer C, Neumann M, Tauschel D, Edelhauser F. Towards non-reductionistic medical anthropology, medical education and practitioner-patient-interaction: The example of anthroposophic medicine, *Patient Educ. Couns*, 89(3), 2012, 455-460.
9. (ATSDR). Toxicological profile for lead (Draft for Public Comment), *ATSDR's Toxicol. Profiles*, 2019, 1-583.
10. Sisman T. Early life stage and genetic toxicity of stannous chloride on zebrafish embryos and adults: Toxic effects of tin on zebrafish, *Environ. Toxicol*, 26(3), 2011, 240-249.
11. Kam P C A, Liew S. Traditional chinese herbal medicine and anaesthesia, *Anaesthesia*, 57(11), 2002, 1083-1089.
12. Ferrando M D, Andreu-Moliner E. Effect of lindane on the blood of a freshwater fish. *Bull. Environ. Contam. Toxicol*, 47(3), 1991, 465-470.
13. Janssen P J M, Bosland M C, Van Hees J P, Spit B J, Willems M I, Frieke Kuper C. Effects of feeding stannous chloride on different parts of the gastrointestinal tract of the rat, *Toxicol. Appl. Pharmacol*, 78(1), 1985, 19-28.
14. Rader J I. Anti-nutritive effects of dietary tin, *In Advances in Experimental Medicine and Biology*, 289, 1991, 509-524.
15. Yousef M I, Awad T I, Elhag F A, Khaled F A. Study of the protective effect of ascorbic acid against the toxicity of stannous chloride on oxidative damage, antioxidant enzymes and biochemical parameters in rabbits, *Toxicology*, 235(3), 2007, 194-202.
16. Yousef M I. Protective role of ascorbic acid to enhance reproductive performance of male

- rabbits treated with stannous chloride, *Toxicology*, 207(1), 2005, 81-89.
17. Beynen A C, Pekelharing H L M, Lemmens A G. High intakes of tin lower iron status in rats, *Biol. Trace Elem. Res*, 35(1), 1992, 85-88.
 18. Yu S, Beynen A C. High tin intake reduces copper status in rats through inhibition of copper absorption, *Br. J. Nutr*, 73(6), 1995, 863-869.
 19. Omura M, Ogata R, Kubo K, Shimasaki Y, Aou S, Oshima Y, Tanaka A, Hirata M, Makita Y, Inoue N. Two-generation reproductive toxicity study of tributyltin chloride in male rats, *Toxicol. Sci*, 64(2), 2001, 224-232.
 20. De Groot A P, Feron V J, Til H P. Short-term toxicity studies on some salts and oxides of tin in rats, *Food Cosmet. Toxicol*, 11(1), 1973, 19-30.

Please cite this article in press as: Mariam A. Ibrahim *et al.* Stannous chloride induces alterations in body weight and hematological parameters in male rabbit, *Asian Journal of Research in Chemistry and Pharmaceutical Sciences*, 9(3), 2021, 92-97.